

KILLS GERMS
IN 30 SECONDS[‡]

FOR DAILY USE

BIOSILVER™

DISINFECTANT, FUNGICIDE & VIRUCIDE*

Disinfects Hospitals, Homes,
Schools, Offices, Restaurants

KILLS
MRSA
& VRE

Disinfects Without Bleaching
No Harsh Chemical Smell

Active Ingredient
Silver† 0.003%
Citric Acid 4.840%
Other Ingredients 95.157%
Total 100.000%

†Electrolytically generated
silver ions

Net Vol. 32 FL OZ

PATENTED
SILVER ION
FORMULA



KEEP OUT OF REACH OF CHILDREN

BIOSILVER™
FAST, EASY, EFFECTIVE

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Biosilver is a colorless, odorless broad spectrum antimicrobial disinfectant and deodorizer. Proven to kill bacteria, fungus and viruses* **Biosilver** should be used on non-porous environmental hard surfaces in hospitals, nursing homes, medical and dental clinics, laboratories, ambulances, funeral homes, hotels, restaurants, schools, day care facilities, nurseries, playrooms, recreational facilities, locker rooms, washrooms, restrooms, offices, veterinary clinics, animal shelters, kennels, exercise facilities, beauty and barber shops, school buses, subways, trains, airplanes, ships and other public transportation vehicles.

Biosilver has been formulated to disinfect hard, non-porous environmental surfaces (painted, glazed tile, plastic, metal, glass, glazed porcelain) and objects including walls, floors, counters, sinks, cabinets, showers, doorknobs, lights switch covers, telephones, bed frames, wheelchairs, examination tables, children's toys, toy boxes, diaper changing tables, play tables, jungle gyms, playhouses, child car seat, strollers, cribs, playpens and activity centers.

STORAGE AND DISPOSAL

Storage: Do not contaminate water, food or feed by storage or disposal. Disposal: Do not reuse container. Rinse thoroughly before discarding in trash or recycling.

IN CASE OF EMERGENCY: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact CHEMTREC 1-800-424-9300 for emergency medical treatment information.



Distributed by:
Mid-Continent Packaging
1200 N 54th St.
Enid, OK 73701

EPA Reg No. 72977-3-66397
EPA Est. No. 66397-OK-1
Made In USA

GENERAL INFORMATION

Biosilver successfully killed the following organisms under AOAC protocols (In order to ensure that all organisms listed are killed, you must use the contact times as directed in the following Application Instructions).

Organism	Kill Time
* <i>Pseudomonas aeruginosa</i>	30 seconds
* <i>Staphylococcus aureus</i>	30 seconds
* <i>Salmonella choleraesuis</i>	30 seconds
* <i>Listeria monocytogenes</i>	30 seconds
Vancomycin resistant <i>Enterococcus faecium</i>	2 minutes
Methicillin resistant <i>Staphylococcus aureus</i>	2 minutes
<i>Escherichia coli</i> 0157:H7	2 minutes
<i>Trichophyton mentagrophytes</i> (Athlete's Foot Fungus)	10 minutes
**HIV type 1- Strain HTLV IIIB	30 seconds
*Herpes Simplex Type 1 VR-733 F(1) Strain	1 minute
*Influenza A VR-544, Hong Kong strain	10 minutes
*Rhinovirus R37 VR-1147, Strain 151-1	10 minutes
*Polio Type 2, VR-1002, Lansing Strain	10 minutes

APPLICATION INSTRUCTIONS

Pre-clean surfaces prior to using this product.

General Disinfection: For general disinfection and control of the bacteria *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Salmonella choleraesuis*, *Listeria monocytogenes*, Vancomycin Resistant *Enterococcus faecium* (VRE), Methicillin Resistant *Staphylococcus aureus* (MRSA) and *Escherichia coli* 0157:H7 the surface must be completely wet with **Biosilver** for 2 minutes. The surface may then be wiped dry with a clean towel. When used as directed, **Biosilver** provides protection from *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella choleraesuis* up to 24 hours after initial application.

Fungus Control:

For effective control of the fungus *Trichophyton mentagrophytes*, the surface must be completely wet with **Biosilver** for 10 minutes. The surface may then be wiped dry with a clean towel. Re-apply when cleaning or when new growth appears.

***Viral Control:** To kill Herpes Simplex Type 1 F(1) Strain, Influenza A Virus, Hong Kong strain, Rhinovirus R37 Strain 151-1, Polio Virus Type 2 Lansing Strain the surface must be completely wet with **Biosilver** for 10 minutes. The surface may then be wiped dry with a clean towel. **Kills HIV-1 on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids** in health care settings (or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV): **Instructions for Cleaning and Decontamination Against HIV** on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids: **Personal Protection:** When handling items soiled with blood or body fluids, use appropriate barrier protection such as latex gloves, gowns, masks or eye coverings. **Cleaning Procedure:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this disinfectant. **Contact Time:** Apply **Biosilver** to area to be treated. The surface must be completely wet with **Biosilver** for 30 seconds. The surface may then be wiped dry with a clean towel. This contact time will not control all organisms listed on this label. Refer to application instructions for other organisms. **Disposal of Infectious Materials:** Blood and other body fluids should be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the human body, either into or in contact with the bloodstream, or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not normally penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.